

THE ROLE OF SEED BANK IN THE DYNAMICS OF UNDERSTORY IN AN OAK FOREST IN HUNGARY

G. KONCZ,^{1,2*} MÁRIA PAPP,¹ P. TÖRÖK,² ZS. KOTROCZÓ,³ ZS. KRAKOMPERGER,²
G. MATUS¹ and B. TÓTHMÉRÉSZ²

¹Department of Botany, University of Debrecen, Debrecen, Hungary

²Department of Ecology, University of Debrecen, Debrecen, Hungary

³Biological Institute, College of Nyíregyháza, Nyíregyháza, Hungary

(Received: June 29, 2010; accepted: August 12, 2010)

We studied the potential role of seed bank in the dynamics of the understory in a turkey-sessile oak forest (*Quercetum petraeae-cerris*) in Hungary. We used long term records of the herb layer (1973–2006) and the seed bank composition of 2006 to assess the role of seed bank in the regeneration of herb layer. The total cover of herb layer decreased from 22% (1973) to 6% (1988), and remained low (<10%) till 2006; coinciding with the increasing cover of secondary canopy dominated by *Acer campestre*. We found a low density seed bank (ca. 1300 seed/m²). Altogether 33 species were germinated from the soil samples. A few generalist weed species composed the majority of seed bank. It was possible to assign a seed bank type for 19 species; 14 species out of 19 was long-term persistent. We found that the characteristic perennial forest herbs and grasses had only sparse seed bank. The Jaccard similarity between vegetation and seed bank was low (<30%). Our results suggest that the continuous establishment of forest herbs are not based on local persistent seed bank; it should be based on vegetative spreading and/or seed rain.

Keywords: Forest decay – herb species – seed density – seed persistence – seedling emergence

INTRODUCTION

The herb layer of forests has quick adaptation features which make it a sensible indicator of forest dynamical changes [1]. It is crucial to assess the composition and density of seed banks to understand dynamical events and predict the responses of the herb layer to disturbance [38, 52, 56, 62]. The seeds in the soil enable the plants to survive the unfavourable conditions for the germination and establishment [6, 23]. Furthermore, the seed bank can determine the direction of secondary succession after a disturbance [44] and can ensure information on the past and also predicts future changes [42]. The research of forest dynamics [17] and forest seed banks [e.g. 4, 27, 41] is in the centre of interest. But only a few studies compared seed bank composition with long term herb layer records [33].

In our study long term vegetation records of the herb layer (1973–2006) were compared with seed bank of 2006 in a turkey oak-sessile oak forest in Hungary. We aimed at to test the following hypotheses: (i) The seed bank is composed by a few

*Corresponding author; e-mail: konczgabo@gmail.com

generalist weed species; the characteristic perennial forest species do not form persistent seed bank [5, 22]. (ii) Species richness and seed density of seed banks are low; only a few short lived species form a persistent seed bank [46, 55]. (iii) Similarity between the forest seed bank and present vegetation is low [48, 51]; the seed bank contains seeds of species characteristic to former composition of vegetation [6]; therefore, the seed bank is more similar to former stages of vegetation than to later ones.

MATERIALS AND METHODS

Site description and history

The study site was in the Síkfökút Project Nature Reserve located ca. 6 km NE of town Eger, in the hilly region of North Hungarian Mountain Range (ca. 47°90'N; 20°46'E, a.s.l. 320–340 m). The climate is moderately continental with a mean annual temperature of 9.9 °C. The mean annual precipitation is between 500–600 mm, the yearly peak of precipitation is in the May–July period. Serious drought events were frequent during the studied period. The 64 ha reserve is vegetated with an even-aged ca. 100 years old xeric oak forest (*Quercetum petraeae-cerris*) [25]. One third of the oak trees perished during the 1980s, and canopy gaps were formed [26, 39, 40]. These gaps were filled up by shrubs and also a dense second canopy was formed by *Acer campestre* [30, 31].

Sampling setup

Cover of vascular herb species were recorded in 33 permanent plots of 4 m × 4 m within a quarter-hectare sized area in July of the following years: 1973, 1982, 1988, 1994, 2000, 2006. Seed bank was sampled after natural winter stratification in the same plots after snowmelt in late March, 2006. In each plot 6 soil cores were collected (4 cm diameter and 10 cm depth, in a total volume of 25,000 cm³). We treated the samples of all plots separately, which were divided in two vertical segments (0–5 cm, 5–10 cm). Identical segments drilled from the same plot were pooled in a bulk reduction procedure used to reduce sample volume [53]. Vegetative organs were retained by washing over a coarse sieve (3.0 mm mesh). Seed-free fine soil components were removed on a 0.2 mm fine mesh. Sample volume was reduced by 65–70%. Volume reduced samples were spread in a 2–4 mm thick layer on 66 trays, formerly filled with 8 cm of steam-sterilized potting soil. Trays were placed under natural light in a greenhouse shaded with Rachel nets from early May to August. Temperature varied typically between 30 °C/18 °C at day/night. Seedlings were regularly counted, identified then removed [10]. Unidentified seedlings were transplanted. In early July, when no new seedlings emerged, watering was stopped; dried sample layers were

crumbled and turned. Early September watering was re-started and continued till late October. Germination altogether lasted for 37 weeks. Seed rain as contamination was monitored in sample-free control trays filled with sterilized soil.

Data management

Raunkiaer life-forms and Grime's C-S-R strategy groups adapted to the Hungarian conditions (Social Behaviour Types) were used for the characterization of herbs [3]. Based on the Social Behaviour Types the followings were regarded as ruderals: AC-alien competitors, DT-disturbance tolerants, NP-natural pioneers, RC-ruderal competitors, W-weeds. We used transient, short term persistent, and long term persistent seed bank type classification of Thompson et al. [56]. Species with less than 3 seeds in the samples (12 seeds/m²) were not classified.

Spearman rank-correlation was used to compare the species ranks of the cover of herbaceous species [49]. Similarity of the seed bank and vegetation data was displayed by nonmetric multidimensional scaling [35]; similarity was calculated by the Jaccard similarity. Differences between years in means of cover and species numbers were tested using RM ANOVA and in case of significant differences Tukey test was used [62].

RESULTS

Understory vegetation

Altogether 53 vascular species were recorded in vegetation; out of these 15 species were recorded only once during the whole study period. The species composition changed substantially during the study period (Table 1). Several characteristic forest herbs (e.g. *Carex michelii*, *C. montana*, *Lathyrus niger*) were typical in the vegetation recording before 1988. From 1988 onwards only sparse cover of forest species were detected (*Lathyrus vernus*, *Melica uniflora*, *Poa nemoralis*). Several nitrophilous were established permanently during the studied period (e.g. *Alliaria petiolata*, *Chelidonium majus*, *Galium aparine* and *Geranium robertianum*). The mean cover of the herb layer was 22% in 1973. This score was decreased below 10% and remained low till the end of the study (Table 2). The mean number of species significantly decreased from 1973 to 1988 ($p < 0.001$), but we found significant increase from 1988 to 2006 ($p < 0.001$, Table 2). We found that the species ranks also changed significantly ($p < 0.001$). The cover of perennial graminoids (e.g. *Poa nemoralis* and *Melica uniflora*, *Carex* species) decreased remarkably. From 1994 onwards the cover of nitrophilous annuals and short-lived ruderals (e.g. *Chelidonium majus*, *Fallopia dumetorum*, *Geranium robertianum*) were increased (Table 1).

Table 1
Percentage cover (mean \pm SE) of the most frequent 25 species during the studied period

Species	SBT	FS	1973	1982	1988	1994	2000	2006
<i>Carex michelii</i>	G	PG	5.9 \pm 1.1	3.4 \pm 0.8	0.3 \pm 0.2			
<i>Carex montana</i>	S	PG	2.8 \pm 0.9	1.4 \pm 0.5				
<i>Dactylis polygama</i>	G	PG	1.7 \pm 0.3	1.0 \pm 0.2				
<i>Fragaria vesca</i>	G	PH	0.4 \pm 0.1	0.2 \pm 0.1				
<i>Festuca heterophylla</i>	C	PG	1.1 \pm 0.4	0.4 \pm 0.1				
<i>Galium schultesii</i>	G	PH	1.5 \pm 0.7	1.2 \pm 0.5	1.2 \pm 0.7			
<i>Lathyrus niger</i>	G	PH	0.5 \pm 0.2	0.4 \pm 0.1				
<i>Melica nutans</i>	G	PH	0.4 \pm 0.2	0.2 \pm 0.1				
<i>Melica uniflora</i>	C	PG	1.4 \pm 0.7	0.8 \pm 0.3		0.5 \pm 0.3	0.3 \pm 0.2	
<i>Poa nemoralis</i>	C	PG	4.6 \pm 1.1	2.5 \pm 0.6	0.5 \pm 0.3	0.2 \pm 0.2	0.1 \pm 0.1	
<i>Alliaria petiolata</i>	DT	PH			0.2 \pm 0.1		0.1 \pm 0.1	
<i>Carex muricata</i>	DT	PG			0.2 \pm 0.1			0.3 \pm 0.1
<i>Galium aparine</i>	W	AH			0.2 \pm 0.1	0.5 \pm 0.3	0.2 \pm 0.1	1.9 \pm 0.8
<i>Geranium robertianum</i>	DT	AH			1.1 \pm 0.8	0.5 \pm 0.2	0.1 \pm 0.1	1.4 \pm 0.9
<i>Lathyrus vernus</i>	S	PH			0.2 \pm 0.1	0.1 \pm 0.1	0.3 \pm 0.1	0.2 \pm 0.1
<i>Fallopia dumetorum</i>	DT	AH			0.5 \pm 0.1	0.8 \pm 0.3	0.3 \pm 0.2	0.2 \pm 0.1
<i>Symphytum nodosum</i>	G	PH			0.2 \pm 0.1			
<i>Chelidonium majus</i>	W	PH				0.8 \pm 0.5	0.1 \pm 0.1	1.3 \pm 0.7
<i>Melittis grandiflora</i>	G	PH				0.1 \pm 0.1		
<i>Polygonatum latifolium</i>	G	PH				0.3 \pm 0.2	0.4 \pm 0.4	0.7 \pm 0.5
<i>Veronica chamaedrys</i>	DT	PH				0.1 \pm 0.1		
<i>Dictamnus albus</i>	S	PH					0.1 \pm 0.1	
<i>Ajuga reptans</i>	DT	PH						0.2 \pm 0.1
<i>Stachys sylvatica</i>	G	PH						0.2 \pm 0.1
<i>Stellaria media</i>	DT	AH						0.2 \pm 0.1

Notations: Social Behaviour Types (SBT, [3]) = C – competitors, DT – disturbance tolerants, G – generalists, W – weeds, S – stress tolerants. Functional species groups (FS) based on morphological categories and Raunkiaer's Life-form system: PG – perennial grass, PH – perennial herb, AH – annual herb

Table 2
Change of vegetation cover and species richness (mean \pm SE) in the study period.
Different superscripted letters refer significant differences between years (RM-ANOVA, $p < 0.001$)

	1973	1982	1988	1994	2000	2006
Cover	22.3 \pm 3.2 ^a	12.2 \pm 1.9 ^b	5.6 \pm 1.3 ^c	4.0 \pm 0.8 ^d	1.8 \pm 0.5 ^e	7.9 \pm 1.4 ^f
Species richness	5.8 \pm 0.4 ^a	5.5 \pm 0.4 ^a	2.4 \pm 0.3 ^b	1.8 \pm 0.3 ^c	1.0 \pm 0.3 ^d	4.2 \pm 0.6 ^e

Table 3

Seed bank composition of the 33 plots in the upper (0–5 cm) and lower (5–10 cm) soil layer (seeds/m²)

Species	SBT	FS	0–5 cm	5–10 cm	P
<i>Ajuga reptans</i>	DT	PH	24	36	3
<i>Ballota nigra</i>	W	PH		4	
<i>Fallopia dumetorum</i>	DT	AH	8	4	2
<i>Carex michelii</i>	G	PG	12		1
<i>Carex muricata</i>	DT	PG	56	68	3
<i>Chenopodium album</i>	RC	AH	52	8	3
<i>Chenopodium hybridum</i>	W	AH	4		
<i>Chenopodium polyspermum</i>	RC	AH	133	193	3
<i>Cherophyllum temulum</i>	DT	AH		4	
<i>Cirsium arvense</i>	RC	PH	4		
<i>Cirsium vulgare</i>	W	PH	32	4	3
<i>Conyza canadensis</i>	AC	AH	48	28	2
<i>Dactylis glomerata</i>	DT	PG	8	4	3
<i>Epilobium montanum</i>	G	PH	8		
<i>Epilobium tetragonum</i>	G	PH	4		
<i>Festuca heterophylla</i>	C	PG	4		
<i>Fragaria vesca</i>	G	PH	24	4	3
<i>Hypericum perforatum</i>	DT	PH	52	52	3
<i>Lathyrus niger</i>	G	PH	4		
<i>Matricaria maritima</i>	W	AH	4		
<i>Melandrium album</i>	W	AH	4		
<i>Melica uniflora</i>	C	PG	4		
<i>Poa nemoralis</i>	C	PG	44	20	2
<i>Polygonum lapathyfolium</i>	DT	PH	4	4	
<i>Solanum nigrum</i>	W	AH	4	8	3
<i>Stellaria media</i>	DT	AH	40	36	3
<i>Taraxacum officinale</i>	RC	PH	4		
<i>Turritis glabra</i>	G	PH	8	4	3
<i>Typha angustifolia</i>	C	PH	52	24	3
<i>Urtica dioica</i>	DT	PH	8		
<i>Verbascum nigrum</i>	G	PH	4	36	3
<i>Veronica chamaedrys</i>	DT	PH	36	8	2
<i>Vicia lathyroides</i>	NP	AH	8	8	3

Abbreviations and notations: the Social Behaviour Types (SBT) categories by Borhidi [3]:=AC – alien competitors, DT – disturbance tolerants, G – generalists, NP – natural pioneers, RC – ruderal competitors, W – weeds. Functional species groups (FS) = PG – perennial grass, PH – perennial herb, AH – annual herb. The seed bank types (P) of species according to the system of Thompson [56]: 1-transient; 2-short-term persistent; 3-long-term persistent.

Seed bank

Altogether 33 species (7 monocots and 26 dicots) emerged from the soil samples. We detected a mean seed density of 1.270 seeds/m². Significantly higher number of seeds were detected in the upper soil layer than in lower one ($p < 0.05$). Altogether 44% of the total seed bank was detected in the lower layer (Table 3). A limited number of species comprised the majority of seed bank. Ten species formed the 80% of total seed bank. *Chenopodium polyspermum* had the highest seed density (ca. 330 seeds/m²). The proportion of ruderals in total seed density was high (about 80%). Out of the perennial graminoids only *Poa nemoralis* (ca. 64 seeds/m²) and *Carex muricata* (ca. 125 seeds/m²) built up detectably dense seed bank. Most of the detected species had at most sparse seed banks (Table 3). Lower than the half of the detected species in the present vegetation (2006) had some seed bank. Altogether for 19 species was possible to assign a seed bank type, and most of them had long-term persistent seeds (14 species).

Herb layer and seed bank

During the whole study period (vegetation 1973–2006 and seed bank in 2006) 158 species were detected. There were 17 species in the seed bank which were not recorded in the vegetation. Among these species *Chenopodium album* and *Typha angustifolia* were the most frequent, but with low mean density (60 and 76 seeds/m², respectively). We found that 37 species of herb layer did not form detectably dense seed




Fig. 1. Nonmetric multidimensional scaling of the species composition of the seed bank and the herbaceous vegetation based on Jaccard similarity (stress = 0.27)

bank. Several among them were characteristic forest species detected frequently in the vegetation (*Melica nutans*, *Polygonatum latifolium*, *Stachys sylvatica*, and *Viola hirta*). Several species (e.g. *Chelidonium majus* and *Geranium robertianum*) detected in vegetation did not form any seed bank (Table 3).

In 2006, 53 species composed the vegetation and the seed bank. Among them 20 species were found exclusively in the vegetation, while 18 species only in the seed bank. There were 15 species both in the vegetation and in the seed bank. The Jaccard similarity between vegetation of the studied years and the seed bank was low (ranged from 11% to 28%; Fig. 1). Higher scores than 20% were only detected between seed banks and the vegetation records of 1973 and 2006, respectively.

DISCUSSION

Density and richness of seed bank

Compared to the literature the detected mean seed density of 1.270 seeds/m² is low. In the broadest sense, in dry-mesophilous grasslands several thousand seeds per square meter is typical [59]; in wet meadows and dry psammophilous grasslands even higher figures are common [37, 38, 57]. In former studies in forest seed banks a wide range of density scores were found. Typical seed density in most of the studies range from several hundred to several thousand seeds per m² in the top 10 cm [11, 24]. Schiffman and Johnson [46] found less than one seed/m² in different stands of Virginian oak forests. In mature beech forests there were found from 1,400 to 21,500 seeds/m² [50] and 12,400 seeds/m² in an oak-beech forest of Belgium [5]. In an old-growth temperate deciduous forest of Quebec there were from 475 to 16,700 seeds/m² [34]. Kjellsson [28] found from 2,200 to 15,600 seeds/m² in Danish deciduous forests. The species richness in the seed bank found in our study, compared to former ones was high (33 species). Godefroid et al. [19] found 9–24 species in temperate forest ecosystem, and only 9–11 species were typical in *Quercus* spp. dominated stands. In mixed coniferous forest Zobel et al. [62] detected 13–29 species.

Species composition of seed banks

We found that 80% of forest seed bank was composed by ruderals. This is well in accordance with the results of previous studies [4, 29, 46]. It was formerly found that, the seed bank of temperate forests was mainly composed by early successional species (characteristic species to clearings), ruderals [4, 5, 22] or rarely by invasive species [22]. For typical species of undisturbed habitats, only short term persistence was proven [42, 58].

The most frequent species was a long-term persistent seeded weed, *Chenopodium polyspermum*. This species is a shade intolerant weed, characteristic to open agricultural areas, and were not detected in the vegetation. As formerly published, the seed

bank density of *Chenopodium* species can exceed 10,000 seeds/m² [16]. Seeds of *Chenopodium polyspermum* – can successfully persist several decades [7, 9, 20, 21, 23, 47]. On the other hand, it is more likely that the detected weed seeds are originated from the surroundings by spatial dispersal. Closely to the studied forest stand abandoned vineyards are situated, where this species is frequently recorded.

Many authors suggest that the most shade-tolerant species do not form a persistent seed bank [e.g. 4, 22]. Our results support these findings; we found that the seed bank of perennial forest grasses and forbs were sparse. The low seed density of these species can be explained with predation, because seeds of late successional species are large and they are more frequently exposed to predation, than small seeded, early successional ones [12, 18]. Moreover, perennial grasses and forbs often could survive strong shading [28]. In such conditions the regeneration and spreading most of the perennial forest species is more likely based on vegetative spreading and reproduction than on seed bank formation [36].

Our third hypothesis was that there is a poor similarity between the present forest vegetation and seed bank composition [54, 60]. As it was presented in the review of Bossuyt and Honnay [2008], that the species similarity between seed bank and vegetation across ecosystems was the lowest in forests with a mean score around 30%. In our study only lower scores were found than that. The main causes of low similarity between vegetation and seed banks of the forests are: (i) Persistent seeds of ruderal species remain viable in the soil for a long time without germination [5, 8, 13, 32]. (ii) In low density seed banks, species with small amounts of short-lived seeds with clumped horizontal pattern can remain undetected by sampling. (iii) Presence of several hygrophytes (e.g. *Typha angustifolia*) exclusively in seed bank is a further source for dissimilarity.

The soil seed bank of the studied oak forest contains many persistent seeds of ruderal species. Conversely, characteristic forest grasses and forbs do not possess dense seed banks. As a consequence, the recovery of former understory typical in the early years of the study (1973–1988) is difficult from local seed banks. There are some other driving forces such as the increased vegetative propagation [14, 15] and seed rain from adjacent areas which have more important influence on the regeneration dynamics of understory in temperate forests than persistent seed bank formation. Our results suggest that the continuous establishment of forest herbs are not based on local persistent seed bank; it should be based on vegetative spreading and/or seed rain.

ACKNOWLEDGEMENT

The authors are thankful to K. Jámbrík, Zs. Veres, P. Ari, A. Miklós, E. Miklós for their help in field sampling and seed germination study. We are thankful to the staff of Debrecen University Botanical Garden for providing facilities as well as L. Karaffa and E. Fekete at Department of Genetics and Applied Microbiology for sterilizing potting soils. Special thanks are due to the workers of Botanical Garden of Debrecen for providing facilities in germination. Research was partially supported by TÁMOP 4.2.1.

REFERENCES

1. Aude, E., Lawesson, J. E. (1998) Vegetation in Danish beech forests: the importance of soil, microclimate and management factors, evaluated by variation partitioning. *Plant Ecol.* 134, 53–65.
2. Bhujju, D. R., Ohsawa, M. (2001) Patch implications in the maintenance of species richness in an isolated forest site. *Biol. Conserv.* 98, 117–125.
3. Borhidi, A. (1995) Social Behaviour Types, the naturalness and relative ecological indicator values of the higher plants in the Hungarian Flora. *Acta Botanica Hungarica* 39, 97–181.
4. Bossuyt, B., Hermy, M. (2001) Influence of land use history on seed banks in European temperate forest ecosystems: a review. *Ecography* 24, 225–238.
5. Bossuyt, B., Heyn, M., Hermy, M. (2002) Seed bank and vegetation composition of forest stands of varying age in central Belgium: consequences for regeneration of ancient forest vegetation. *Plant Ecol.* 162, 33–48.
6. Bossuyt, B., Honnay, O. (2008) Can the seed bank be used for ecological restoration? An overview of seed bank characteristics in European communities. *J. Veg. Sci.* 19, 875–884.
7. Canham, C. D., Marks, P. L. (1985) The response of woody plants to disturbance, patterns of establishment and growth. In: Pickett, S. T. A., White, P. S. (eds.) *The Ecology of Natural Disturbance and Patch Dynamics*. Academic Press, New York, pp. 197–216.
8. Chang, E. R., Jefferies, R. L., Carleton, T. J. (2001) Relationship between vegetation and soil seed banks in an arctic coastal marsh. *J. Ecol.* 89, 367–384.
9. Cook, R. (1980) The biology of seeds in the soil. In: Solbrig, O. T. (ed.) *Demography and evolution of plant populations*. University of California Press, Berkeley, pp. 107–129.
10. Csapody, V. (1968) *Keimlingsbestimmungsbuch der Dicotyledonen*. Akadémiai Kiadó, Budapest.
11. Csontos, P. (2007) Seed banks: ecological definitions and sampling considerations. *Community Ecol.* 8(1), 75–85.
12. Decocq, G., Valentin, B., Toussaint, B., Hendoux, F., Saguez, R., Bardat, J. (2004) Soil seed bank composition and diversity in a managed temperate deciduous forest. *Biodivers. Conserv.* 13, 2485–2509.
13. Drake, D. R. (1998) Relationships among the seed rain, seed bank and vegetation of a Hawaiian forest. *J. Veg. Sci.* 9, 815–828.
14. Eriksson, O. (1989) Seedling dynamics and life histories in clonal plants. *Oikos* 55, 231–238.
15. Eriksson, O. (1993) Dynamics of genets in clonal plants. *Trends Ecol. Evol.* 8, 313–316.
16. Forester, E. (1956) Ein Beitrag zur Kenntnis der Selbstverjüngung von Dauerweiden. *Zeitschrift für Acker- und Pflanzenbau* 100, 273–301.
17. Frelich, L. E. (2002) *Forest Dynamics and Disturbance Regimes: Studies from Temperate Evergreen-Deciduous Forests*. Cambridge University Press, Cambridge.
18. Gashaw, M., Michelsen, A., Jensen, M., Friis, I. (2002) Soil seed banks dynamics of fire-prone wooded grassland, woodland and dry forest ecosystems in Ethiopia. *Nord. J. Bot.* 22, 5–17.
19. Godefroid, S., Phartyal, S. S., Koedam, N. (2006) Depth distribution and composition of seed banks under different tree layers in a managed temperate forest ecosystem. *Acta Oecol.* 29, 283–297.
20. Grime, J. P. (1989) Seed banks in ecological perspective. In: Leck, M. A., Parker, V. T., Simpson, R. L. (eds) *Ecology of Soil Seed Banks*. Academic Press, Inc., London, pp. 15–22.
21. Grubb, P. J. (1988) The uncoupling of disturbance and recruitment, two kinds of seed bank and persistence of plant populations at the regional and local scales. *Ann. Zool. Fenn.* 25, 23–26.
22. Halpern, C. B., Evans, S. A., Nielson, S. (1999) Soil seed banks in young closed-canopy forests of the Olympic Peninsula, Washington: potential contributions to understory reinitiation. *Can. J. Bot.* 77, 922–935.
23. Harper, J. L. (1977) *Population biology of plants*. London, Academic Press.
24. Hopfensperger, K. (2007) A review of similarity between seed bank and standing vegetation across ecosystems. *Oikos* 116, 1438–1448.
25. Jakucs, P. (1985) *Results of "Sikfőkút Project"*. Akadémiai Kiadó, Budapest.
26. Jakucs, P., Mészáros, I., Papp, B. L., Tóth, J. A. (1986) Acidification of soil and decay of sessile oak in the "Sikfőkút Project" area (N-Hungary). *Acta Bot. Hung.* 32, 303–322.

27. Kalamees, R., Zobel, M. (1998) Soil seed bank composition in different successional stages of a species rich wooded meadow in Laelatu, western Estonia. *Acta Oecol.* 19, 175–180.
28. Kawano, S. (1975) The productive and reproductive Biology of Flowering plants. II. The concept of Life History Strategy in plants. *J. College of Libertal Arts* 8, 51–86.
29. Kjellsson, G. (1992) Seed bank sin Danish deciduous forests: species composition, seed influx and distribution pattern in soil. *Ecography* 15, 86–100.
30. Kotroczó, Zs., Fekete, I., Tóth, J. A., Tóthmérész, B., Balázsy, S. (2008) Effect of leaf- and root-litter manipulation for carbon-dioxide efflux in forest soil. *Cereal Res. Commun.* 36 (Supl.), 663–666.
31. Krakomperger, Zs., Tóth, J. A., Varga, Cs., Tóthmérész, B. (2008) The effect of litter input on soil enzyme activity in an oak forest.: *Cereal Res. Commun.* 36 (Supl.), 322–326.
32. Leary, C. I., Howes-Keiffer, C. (2004) Comparison of standing vegetation and seed bank composition one year following hardwood reforestation in south-western Ohio. *Ohio J. Sci.* 104, 20–28.
33. Leck, C. F. (1998) A ten-year seed bank study of old field succession in central New Jersey. *J. Torrey Bot. Soc.* 125, 11–32.
34. Leckie, S., Vellend, M., Bell, G., Waterway, M. J., Lechowicz, M. J. (2000) The seed bank in an oldgrowth, temperate deciduous forest. *Can. J. Bot.* 78, 181–192.
35. Legendre, P., Legendre, L. (1998) *Numerical Ecology*. Elsevier Science, Amsterdam, The Netherlands.
36. Matlaga, D. P., Horvitz, C. C. (2009) Growth and survival across a gap-understory gradient: contrast in performance of sexually vs. clonally produced offspring. *Am. J. Bot.* 96, 439–447.
37. Matus, G., Tóthmérész, B., Papp, M. (2003) Restoration prospects of abandoned species-rich sandy grassland in Hungary. *Appl. Veg. Sci.* 6, 169–178.
38. Matus, G., Papp, M., Tóthmérész, B. (2005) Impact of management change on vegetation dynamics and seed bank formation of inland dune grassland in Hungary. *Flora* 200, 296–306.
39. Mészáros, I., Módy, I., Marschall, M. (1993) Effect of air pollution on the condition of sessile oak forests in Hungary. In: Vernet, J.-P. (ed.) *Environmental Contamination*. Studies in Environmental Science 55, Elsevier Sci. Publ., Amsterdam, pp. 23–35.
40. Mészáros, I., Veres, Sz., Kanalas, P., Oláh, V., Szöllősi, E., Sárvári, É., Lévai, É., Lakatos, Gy. (2007) Leaf growth and photosynthetic performance of two co-existing oak species in contrasting growing seasons. *Acta Silv. Lign. Hung.* 3, 7–20.
41. Milberg, P. (1995) Soil seed bank after eighteen years of succession from grassland to forest. *Oikos* 72, 3–13.
42. Mitlacher, K., Poschlod, P., Rosén, E., Bakker, J. P. (2002) Restoration of wooded meadows – a comparative analysis along a chronosequence on Öland Sweden. *J. Veg. Sci.* 5, 63–73.
43. Numata, M. (1979) Facts, causal analyses, and theoretical considerations on plant succession. *Vegetation und Landschaft Japans* 16, 71–91.
44. Pakeman, R. J., Small, J. L. (2005) The role of the seed bank, seed rain and the timing of disturbance in gap generation. *J. Veg. Sci.* 16, 121–130.
45. Peterson, C. J., Carson, W. P. (1996) Generalizing forest regeneration models: the dependence of propagule availability on disturbance history and stand size. *Can. J. Forest Res.* 26, 45–52.
46. Pickett, S. T. A., McDonnell, M. J. (1989) Seed bank dynamics in temperate deciduous forest. In: Leck, M. A., Parker, V. T., Simpson, R. L. (eds) *Ecology of Soil Seed Banks*. Academic Press, Inc., London, pp. 123–145.
47. Priestly, D. A. (1986) *Seed aging: implications for seed storage and persistence in the soil*. Cornell University Press, Ithaca, N.Y.
48. Schiffman P., Johnson, W. C. (1992) Sparse buried seed bank in a Southern Appalachian oak forest: Implication for succession. *Am. Midl. Nat.* 127, 258–267.
49. Sokal, R. R., Rohlf, F. J. (1995) *Biometry*. Freeman, New York, USA.
50. Staaf, H., Jonsson, M., Olsen, L. G. (1987) Buried germinative seeds in mature beech forests with different herbaceous vegetation and soil types. *Holarctic Ecology* 10, 268–277.
51. Sullivan, K. A., Ellison, A. M. (2006) The seed bank of hemlock forests: implications for forest regeneration following hemlock decline. *J. Torrey Bot. Soc.* 133, 393–402.

52. Templeton, A. R., Levin, D. A. (1979) Evolutionary consequence of seed pools. *Am. Nat.* 114, 232–249.
53. ter Heerdt, G. N. J., Verweij, G. L., Bekker, R. M., Bakker, J. P. (1996) An improved method for seed bank analysis: seedling emergence after removing the soil by sieving. *Funct. Ecol.* 10, 144–151.
54. Thompson, K., Grime, J. P. (1979) Seasonal variation in the seed banks of herbaceous species in ten contrasting habitats. *J. Ecol.* 67, 893–921.
55. Thompson, K. (1992) The functional ecology of seed banks. In: Fenner, M. (ed.) *Seeds: the ecology of regeneration in plant communities*. CAB International, Wallingford, UK, pp. 231–258.
56. Thompson, K., Bakker, J. P., Bekker, R. M. (1997) *The soil seed banks of North West Europe: methodology, density and longevity*. Cambridge University Press, Cambridge, UK.
57. Török, P., Deák, B., Vida, E., Valkó, O., Lengyel, S., Tóthmérész, B. (2010) Restoring grassland biodiversity: Sowing low-diversity seed mixtures can lead to rapid favourable changes. *Biol. Conserv.* 143, 806–812.
58. Turner, M. G. (1989) Landscape ecology: the effect of pattern on process. *Ann. Rev. Ecol. Evol. S.* 20, 171–197.
59. Valkó, O., Török, P., Matus, G., Tóthmérész, B. (2010) Restoration potential in seed banks of acidic fen and dry-mesophilous meadows: can restoration be based on local seed banks? *Restor. Ecol.* doi: 10.1111/j.1526-100X.2010.00679.x
60. Warr, J. S., Kent, M., Thompson, K. (1994) Seed bank composition and variability in five woodlands in southwest England. *J. Biogeog.* 21, 151–168.
61. Zar, J. H. (1999) *Biostatistical Analysis*. Prentice & Hall, Upper Saddle River, New Jersey.
62. Zobel, M., Kalamees, R., Püssa, K., Roosalu, E., Moora, M. (2007) Soil seed bank and vegetation in mixed coniferous forest stands with different disturbance regimes. *Forest Ecol. Manag.* 250, 71–76.